

CrossRef DOI of original article:

1 Ogluzdin V.E.

2 Received: 1 January 1970 Accepted: 1 January 1970 Published: 1 January 1970

3

4 **Abstract**

5 What happens to photons of laser radiation whose energy differs from the energy of the
6 inter-level electronic transitions of atoms of the substance under study. According to Bohr and
7 Einstein, the traditional scheme of interaction between radiation and matter includes the
8 following processes: 1) resonant absorption of radiation, 2) spontaneous radiation, 3) resonant
9 stimulated radiation. In the absence of resonance between the pumping frequency and the
10 frequencies of electronic transitions in atoms alloying a dispersed medium, the interaction of
11 radiation and matter is carried out due to the annihilation of pairs of pump radiation photons,
12 the birth of axions and their subsequent decay into new photons whose energy differs from the
13 energy of the pump photons. If in atomic vapors in the resonant case one photon is required
14 for the transition of an electron from the ground level S to the excited P, then in the
15 non-resonant case such a transition is possible due to the annihilation of a pair of photons.

16

17 **Index terms—**

18 **1 Introduction**

19 hat happens to photons of laser radiation whose energy differs from the energy of the inter-level electronic
20 transitions of atoms of the substance under study. According to Bohr and Einstein, the traditional scheme of
21 interaction between radiation and matter includes the following processes: 1) resonant absorption of radiation,
22 2) spontaneous radiation, 3) resonant stimulated radiation. In the absence of resonance between the pumping
23 frequency and the frequencies of electronic transitions in atoms alloying a dispersed medium, the interaction of
24 radiation and matter is carried out due to the annihilation of pairs of pump radiation photons, the birth of axions
25 and their subsequent decay into new photons whose energy differs from the energy of the pump photons. If in
26 atomic vapors in the resonant case one photon is required for the transition of an electron from the ground level
27 S to the excited P, then in the non-resonant case such a transition is possible due to the annihilation of a pair of
28 photons.

29 **2 II.**

30 **3 Axions in A Dispersed Medium**

31 First of all, we should recall the definition of a dispersed medium (DM). In DM, the doping (alloying) phase in
32 the form of small formations is distributed in the main volume of the continuous phase. Continuous phase DM
33 -glass, liquid, air, vacuum. The second is the doping component DM, weighted in the volume of the continuous
34 phase: these are atoms, molecules or nanoparticles of an element alloying the medium. Air, water, organic
35 liquids, atomic vapors of alkali metals in a vacuum cuvette are also DM. When a laser beam propagates in DM,
36 its spectral and angular characteristics change.

37 The aim of the work is to analyze the processes accompanying the propagation of a laser beam in DM in
38 the absence of resonance between the pumping frequency and the frequencies of electronic transitions. The high
39 density of photons in the laser beam and the high intensity of the electromagnetic field of the atomic nuclei of the
40 element selected for DM doping cause the possibility of annihilation of photon pairs, which leads to the birth of
41 axions. Their decay determines the features of the spectral and angular structures of radiation at the exit from
42 the medium.

43 What happens to the photons of laser radiation in DM? What do we have at the output of DM? How does
44 the radiation spectrum change?

4 A) FORCED ELECTRON RAMAN SCATTERING (EFRS) IN ATOMIC POTASSIUM VAPORS

45 Only in a vacuum does the propagation of light radiation obey the laws of wave optics. In this case, laminar
46 collisionless movement of photons in the beam occurs. In the case of propagation of light radiation in the DM,
47 due to the interaction of the photons of the beam with the electron shell of the atomic nuclei of the atoms alloying
48 the medium and the atoms of the medium itself, the laminar motion of the photons is disrupted, the velocity of
49 the photons meeting the electron shell of the atoms alloying the medium slows down. This circumstance leads to
50 interphoton collisions, interphoton interaction, annihilation of photon pairs with the birth of axions in the strong
51 field of the atomic nucleus [1].

52 According to the works [2] of the American physicist Primakov (HENRY PRIMAKOFF), annihilation (fusion)
53 of two quanta (photons) in the electro-magnetic field of the atomic nucleus can lead to the birth of the axion -A?
54 Leaving the atom, the axion decays into two new quanta (photons) -the forward and reverse Primakov effects:
55 $h? + h? = A? = h? i j + h? 0 j$ (1)

56 where: $h?$ is the energy of quanta (photons) of light radiation used to pump $D?$, $?$ is the frequency of this
57 radiation. Just note that the Primakov effect may be direct or reverse. In the second case, when you turn off
58 the field of the atomic nucleus decays of the axion to two photons. $h? i j$ -is the energy of quanta (photons) of
59 radiation from one of the many components of the broadened spectrum of radiation scattered by the medium, or
60 one of the many components of the photoluminescence spectrum, or one of the many components of the angular
61 spectrum of radiation at the output of the investigated $D?$, (determined by the difference between the energy of
62 the virtual level $-i$ and the energy of atomic level j ; The energy of the virtual level is equal to the sum of the
63 energies of two pump photons. The energy of the virtual level $-i$ can be greater or less than the energy of the
64 electronic levels of the doping element under study. $h? 0j$ -is the energy of the quanta absorbed by the medium;
65 the energy of this quantum is equal to the energy of the electron transition from the ground level to the excited
66 one .The number of levels j in the spectrum of the atom of the element under study is infinite.

67 What determines the energy of the virtual level? The energy of the virtual level -can be greater or less than
68 the energy of the electronic levels of the alloying element under study.

69 Since the birth of the axion and its existence presupposes the presence of strong electromagnetic fields, in our
70 case, the fields of the atomic nucleus, neither Rayleigh scattering on solid particles nor Mandelstam-Brillouin
71 scattering by condensed media (solids and liquids) as a result of its interaction with the intrinsic elastic vibrations
72 of these media will be considered by us. We will also not take into account the scattering of light by fluctuations
73 in the density of the medium (small local deviations of the density from its average value), with which, as is
74 commonly believed in the wave model, the scattering process is associated.

75 4 a) Forced electron raman scattering (EFRS) in atomic potassium vapors

76 It was experimentally possible to register the process of annihilation of photonic pairs and the decay of axions
77 in the case of almost resonant interaction of laser radiation with atomic potassium vapor in the frequency range,
78 where the normal dispersion and where the refractive index is greater than one. These conditions in atomic
79 potassium correspond to the electronic transition $4S 1?2 -4P 3?2$ (Fig. 1). We will consider a region of the
80 spectrum where interphoton interaction is possible, annihilation of photon pairs, the birth of axions and their
81 decay are possible.

82 As a rule, the energy of laser-pumped photons does not coincide with the inter-level energy interval of any
83 of the electronic transitions of both the alloying atom and the atoms and molecules of the doped medium. A
84 frequency-tunable parametric light generator (PLG) was used as a pumping laser in [3], which made it possible
85 to study changes in the frequency and angular structure of the spectrum at the output of the cuvette in atomic
86 pairs of cadmium in the frequency range of the main doublet. The radiation power of PLG was 50...500 kW
87 with a pulse duration of 10...15 ns. The temperature of potassium vapors in the cuvette was maintained in the
88 range of 250...300°C. The radiation spectrum at the outlet of the cuvette was recorded on a spectrograph with a
89 dispersion of 6 A/mm . Recall the frequencies corresponding to potassium doublet lines: lowfrequency doublet
90 line $4S 1?2 -4P 1?2 : ? 02 = 12989 \text{ cm}^{-1}$, high-frequency doublet line $4S 1?2 -4P 3?2 : v 01 = 13046 \text{ cm}^{-1}$.
91 This experiment is described in detail in [3][4][5].

92 In the first spectrogram (Fig. 1a), the pump radiation frequency is $v ? 13020 \text{ cm}^{-1}$, less than the transition
93 frequency $4S 1?2 -4P 3?2$. In this case, with a shift to the low-frequency region of the spectrum relative to the
94 pumping frequency $-?$, a resonant electron forced raman scattering (EFRS) line is recorded at the output of the
95 cuvette, the frequency of which is indicated by $? 3 : ? 3 = 2 ? -? 01$.

96 (2)

97 Calculated according to the ratio (2), the frequency value $? 3$ coincides with the experimental value.

98 The energy of photons of radiation at the frequency $? 3$ can be obtained from the ratio:
99 $h? 3 = 2h? -h? 01$

100 .(3)

101 In relation (3) there is a term $2h?$, indicating the addition of two pump radiation photons (annihilation of
102 which in the field of the atomic nucleus leads to the appearance of an axion -the Primakov effect). In the field of
103 the same atomic nucleus, in an elementary act, an axion whose energy is equal to $2h?$, decays into two photons
104 (quanta) -the reverse Primakov effect.

105 Laser radiation (frequency $v ? 13020 \text{ cm}^{-1}$) transfers electrons from the ground level $4S 1?2$ to the virtual

106 one, the energy of which is determined by the value $E = 2 \hbar$? with the formation of an axion. The energy of the
107 virtual level in the atom is counted from the main level. When the axion decays, the electron transitions from
108 the virtual level $2 \hbar$? to the level $4P 3?2$.

109 The radiation line of the pump? at the outlet of the cuvette is widened. The question of broadening the laser
110 radiation lines in a dispersed medium was considered in [1]. The broadening of the lines is associated with the
111 annihilation of photon pairs, the appearance of axions and their decay. Accordingly, with the pump? radiation
112 line at the output of the cuvette, the EFRS? 3 line is also widened, which follows from the ratio (2).

113 In the spectrogram (Fig. 1a), in addition to the pump? radiation and the radiation at the frequency of 3 ,
114 there is a complex spectral structure in the antistokes region of the spectrum relative to the pump (from $?$ to? as
115), reflecting the change in the divergence (angular spectrum) of radiation at the output of the cuvette.

116 The features of the spectrum associated with the change in the angular spectrum of radiation at the output
117 of the cuvette at the frequencies $?, ? 3 , ?$ as , we will consider in the next section. $3 2? -v 01$ (the value of
118 $2?$ reflects the process of annihilation of two photons with the formation of an axion, during the decay of which
119 photons appear at the frequency 3). The reference lines of the doublet $v 02 , v 01$ on the spectrogram 1 b) were
120 obtained using a spectral lamp. In the second spectrogram (Fig. 1b), the pump radiation frequency 13070 cm^{-1} is shifted to the anti-Stokes region relative to the frequency of the transition line $4S 1?2 -4P 3?2$. In this case,
121 we have a broadening of the frequency spectrum. On a spectrogram with a shift to the high-frequency region
122 of the spectrum relative to the pumping frequency $?$, a weak line of resonant electron forced raman scattering
123 (EFRS) is also recorded at the output of the cuvette, the frequency of which is indicated by 3 . Thus, the
124 EFRS process observed at small detunings of the pump frequency from the resonant transition frequency in
125 potassium vapor during the absorption of two photons in an elementary act with the formation of an axion is
126 accompanied by the transfer of electrons from the main level $4S 1?2$ to the level $4P 3?2$. What distinguishes
127 an almost resonant process from a resonant one. In the case of resonant pumping, the transfer of an electron
128 from the $4S 1?2$ level to the $4P 3?2$ level is carried out due to single-photon absorption. b) The structure of the
129 angular spectrum of the laser beam, behind the output window of the cuvette with atomic potassium vapor When
130 considering the angular spectrum of the laser beam at the output of the cuvette, in the frequency range close to
131 the frequency of the resonant transition, self-action processes (self-focusing, self-focusing) take place [3,5], which
132 do not imply a change in the structure of the frequency spectrum of laser radiation at the output of the medium.
133 The results of the observation of these processes are presented in [6,7]. The effects of self-action are associated
134 with an uneven, Gaussian distribution of the radiation intensity over the crosssection of the laser beam. Near
135 the frequency of the resonant transition, the refractive index of the medium (in our case of atomic potassium
136 vapors) in the electromagnetic field of the laser beam depends on the intensity distribution over the cross-section
137 of the beam. This circumstance explains the change in the wavefront of the beam, leading either to self-focusing
138 (01) or to self-focusing (01).

140 Let us turn once again to the spectrograms presented in Fig. 1. The optical scheme of their registration is
141 constructed in such a way that frequencies are recorded on the spectrograms outside the output window of the
142 cell in the horizontal direction relative to the axis of the laser beam, and in the vertical direction the radiation
143 propagating at an angle to the axis of the laser beam is recorded. To do this, a lens was installed between the
144 output window of the cuvette and the slit of the spectrograph, displaying the output window of the cuvette on
145 the slit of the spectrograph. The slit displays the diameter of the base of the cone of light radiation resting on
146 the exit window of the cuvette. Such an optical scheme makes it possible to register the frequency spectrum of
147 the cone components of radiation scattered in potassium vapor at an angle to the optical axis.

148 Consider the features of the angular spectrum of radiation at the output of the cuvette in the case when the
149 radiation frequency of the pump is v ? $v 01$. In this case, the structure of the angular spectrum at the frequencies
150 v , $v 3$, as well as in the frequency range between v and v as is recorded on the spectrogram.

151 First of all, we note (Fig. 1a) that at the frequency v outside the output window of the cell, the central spot
152 corresponds to the initial direction of propagation of the laser beam. The radiation shifted up and down relative
153 to the optical axis of the frequency v at the same angle indicates that a divergent cone of radiation is formed in
154 the cuvette, the base of which is the output window of the cuvette.

155 In the considered region of the spectrum, the refractive index is greater than one. In such an environment, the
156 speed of photons outside the laser beam is equal to the phase speed of light at the frequency v ? c ? n . When a
157 laser beam propagates in atomic potassium vapors, first of all, the populations of the $4S$ and $4P$ levels align on
158 the beam axis, which changes the refractive index. Under these conditions, it is equal to one. Therefore, along
159 the beam axis, both photons and axions can propagate at the speed of light $?$. According to the definition, the
160 propagation of particles (in our case axions) at the speed of light in a medium where photons move at the phase
161 speed of light is accompanied by Vavilov-Cherenkov radiation. Two spots shifted up and down at the same angle
162 indicate the birth of such radiation in a cuvette with potassium vapors.

163 Let's return to Fig. 1a. At the frequency $v 3$ we have a similar picture -a central spot and components scattered
164 at an angle up and down. According to relation (2), their appearance is caused by the process of three-photon
165 electron scattering, which repeats the structure of the radiation spectrum at the frequency v for the frequency v
166 3 .

167 Finally, we need to explain the structure of the angular spectrum in the frequency range from v to v_{as} . In [8]
168 this structure was called "mustache". It is interesting because there are no pump photons on the beam axis near

7 CONCLUSION

169 the frequency ν as . In [6], it was proposed to use nonlinear polarization to interpret this structure.. The absence
170 of pump radiation photons on the optical axis does not exclude the presence of axions there -particles propagating
171 in the medium at the speed of light and responsible for the Vavilov-Cherenkov radiation cone. This interpretation
172 of the nature of the "mustache" strictly corresponds to the definition of Vavilov-Cherenkov radiation, according
173 to which it is the beam of particles (axions), moving at the speed of light that, causes this radiation.

174 5 c) Propagation of laser radiation in transparent media

175 (air, water); heating of the medium In the case of propagation of laser beams in transparent media (air, water),
176 light scattering occurs, which can be explained by the decay of axions in the electromagnetic fields of nitrogen,
177 oxygen and hydrogen atoms. Let us analyze the case of the propagation of a light beam in a medium, considered
178 in [9]. "Let's assume that we illuminate a transparent medium in complete darkness, for example, clean water,
179 with an intense laser beam. Even if the medium does not contain any impurity particles, the trajectory of the
180 beam in the medium may become slightly noticeable even when observed in directions that do not lie in the plane
181 of incidence?We must... reveal... the origin of this weakly scattered light in all directions, which is superimposed
182 on a more intense unidirectional beam." The authors [9] believe that fluctuations in the density of the medium
183 (in this case, water) are responsible for the scattering of light in all directions. However, this explanation is not
184 enough, part of the energy of the laser beam is spent on heating water. Scattering by density fluctuations does
185 not lead to heating of water.

186 In our opinion, it is natural to assume that when axions decay into two photons at a frequency ν_{ij} , radiation
187 scattering occurs in all directions (angle 4π steradian). It is possible to explain the heating of the medium,
188 which is associated with the process of nonradiative relaxation to the ground level of electrons trapped at the
189 excited level when the medium absorbs the second quantum of the decayed axion in accordance with the ratio
190 (1).It can also be assumed that similar processes occur in the earth's atmosphere under the influence of solar
191 radiation; solar photons annihilate in the earth's atmosphere with the formation of axions, the decay of which is
192 associated with the heating of the atmosphere and possibly photons at frequencies ν_{ij} contribute to the blue
193 color of the sky.

194 Let's add a few remarks to the question of what role axions play in laser technology. We found out that in the
195 absence of resonance between the pump radiation frequency and electronic transitions in the medium, the axion
196 decay involves the transfer of thermal energy to the medium and its heating due to the process of non-radiative
197 relaxation. Therefore, cooling systems are used for heat removal in solid-state lasers, and active medium pumping
198 systems are used in dye lasers.

199 The process of scattering radiation in all directions (angle 4π steradian) in [10] was used to illuminate the
200 laser mirror with the reversal of the wave front. Such a laser works in the presence of phase in homogeneities in
201 the resonator.

202 6 III.

203 7 Conclusion

204 According to Bohr and Einstein, the traditional scheme of interaction between radiation and matter includes
205 the following processes: 1) resonant absorption of radiation, 2) spontaneous radiation, 3) resonant stimulated
206 radiation. In the absence of resonance between the pumping frequency and the frequencies of electronic transitions
207 in atoms alloying a dispersed medium, the interaction of radiation and matter is carried out due to the annihilation
208 of pairs of pump radiation photons, the birth of axions and their subsequent decay into new photons whose energy
209 differs from the energy of the pump photons. Let's list the processes for which the use of the axiope model seems
210 natural. This is the scattering of light in dispersed media, including scattering in air by a solid angle equal to
211 4π steradian. This is resonant forced raman scattering of electrons (EFRS). The decay of the axion into two
212 quanta, whose energy differs from the energy of the pump radiation photons, explains the heating of the medium
213 by the incident pump radiation. This is also the photoluminescence process considered in [11].

214 Finally, an axion is a particle generating a cone of Cherenkov-Vavilov radiation under specially created
215 experimental conditions.



1

Figure 1: Fig. 1 :

216 [Yu et al. ()] , M Yu , S G Kirin , A M Rautian , B M Semenov , Chernobrod . *Pis'maZh. Éksp. Teor. Fiz* 1970.
217 11 p. 340.

218 [Akhmanov et al. ()] , S A Akhmanov , A I Kovrigin , S A Maksimov , V E Ogluzdin . *Pis'maZh. Éksp. Teor.*
219 *Fiz* 1972. 15 p. .

220 [Bagan et al. ()] , A A Bagan , V B Gerasimov , A V Golyanov , V E Ogluzdin , Kvatovaya Electronika . 1990.
221 1 p. .

222 [Bohren and Huffman ()] *Absorption and Scattering of Light by Small Particles*, C F Bohren , D R Huffman .
223 1983. New York?Chichester?Brisbane?Toronto; Singapore: John Wiley & Sons Inc.

224 [Ogluzdin and Vavilov ()] 'Cherenkov effect under condition of near resonant interaction of intense light beams
225 with atomic potassium vapor'. V E Ogluzdin , Vavilov . *Zh.Eksp. Teor. Phyz* 1980. 79 p. .

226 [Ogluzdin ()] *Investigation of the processes of propagation of coherent radiation pulses in resonant media with
227 inverted and non-inverted populations. PhD thesis It is protected in the council in K-053 the specialty
228 Kvantovaja Electron at the Moscow State University named after M, V E Ogluzdin . 1980. p. .*

229 [Primakoff ()] 'Photo-Production of Neutral Mesons in Nuclear Electric Fields and the Mean Life of the Neutral
230 Mesons'. H Primakoff . *Phyz. Rev* 1951. 81 p. .

231 [Ogluzdin ()] 'Photons traveling at the speed of light in two-level atomic medium as source of Cherenkov radiation
232 cones'. V E Ogluzdin . *Physics-Uspekhi* 2004. 47 p. .

233 [Anikin et al. ()] *Resonant electron forcedraman scattering in potassium vapor. Dispersion near the main doublet
234 and the influence of four-photon processes*, V I Anikin , S V Kryuchkov , V E Ogluzdin . 1974. 1991-1998.
235 *Kvantovaja Electron*. 1.

236 [Ogluzdin and The ()] 'Role of Axions in the Formation of the Photoluminescence Spectrum in Dispersive Media'.
237 V E Ogluzdin , The . 10.3390/foundations2010011. <https://doi.org/10.3390/foundations2010011>
238 Foundations 2022. 2 p. .

239 [Ogluzdin (2023)] 'The Role of [Axions in the Formation of Frequency and Angular Spectra of Laser Radiation
240 Propagated in Dispersion'. V E Ogluzdin . *Medium Journal of Applied Spectroscopy* 2023. January. November-
241 December). 89 (6) . (Russian Original 2022)